

Mode of action

Systemic fungicide with protective, curative and eradicant action.

Use/ Application

Crops: **Asparagus** (Rust, powdery mildew, Stemphylium); **Bean and groundnut** (Rust);

Crucifers (Alternaria); **Cucurbits** (Powdery mildew) **Garlic and Onion**

Alternaria, Botrytis, Stemphylium); **Groundnut** (Cercospora, rust); **Pea**

(Powdery mildew); **Banana** Mycosphaerella); **Grapes** (Botrytis, Powdery

mildew); **Mango and papaya** (Powdery mildew); **Orchids** (Fungal leaf spots)

Ornamentals (Powdery mildew, rust, botrytis)

Dosage:

Per hectare (600 L Water)	Per Arpent (250 L Water)	Per Litre of Water
600 ml	250 ml	1ml

Mixing instructions: Prior to pouring, half fill the tank with water and apply agitation while adding the required amount of the product to the spray tank.

Number of applications: Do not use more than four times per crop cycle.

Warning

Re-entry period: 24 hrs

Use restrictions: Avoid repeated use during a crop cycle.

Compatibility: Incompatible with cuprox, copper oxychloride, champion,

funguran - OH, kocide 101

AGRILIX 25% EW

Tebuconazole (a.i) 250g/L

Fungicide

Triazole

POISON

W.H.O CLASS: III

Acute oral LD50(rat):>4264 mg/kg

Acute Dermal LD50(rat):>2000 mg/kg

Note to physician: No specific antidote,

Symptomatic treatment

Manufacturer/ Distributor

Willowood Ltd



Importer/Distributor:

The Mauritius Co-operative Agricultural Fed. Ltd



United Docks Business Park Caudan, Port Louis

Tel: 211 0928/212 1360 Fax: 211 2261

Date of Manufacture:

Batch No.:

DCCB Registration No.:AC/09/06/120

Toxicity: Non toxic to bees, fish; Toxic to animals.

Inhalation toxicity: LC50: 2.5 mg/l/Rat

Pre-harvest interval (days): Garlic, onion (7); Asparagus, bean, crucifers, cucurbits, pea, banana, groundnut, grapes, mango and papaya (14)

Precautions

- Wear protective equipment(overall,gloves, goggles, respirator and boots) when handling and applying product
- Do not eat, drink or smoke while handling pesticide
- Avoid inhalation of spray mist
- Remove contaminated clothing immediately
- After pesticide handling, immediately wash hands and face thoroughly with soap and water
- Keep out of reach of children and store under lock and key
- Do not contaminate ponds and rivers
- Do not re-use empty containers
- Destroy and bury empty containers
- In case of poisoning, seek medical advice
- Storage: Store away from seeds, fertilisers and compost.

QUANTITY: 500 ml

